

Evolution of differential maternal age effects on male and female offspring development and longevity

Martin I. Lind^{*,†,1}, Elena C. Berg^{*,†,1,2}, Ghazal Alavioon¹ and Alexei A. Maklakov¹

¹Ageing Research Group, Department of Animal Ecology, Evolutionary Biology Centre, Uppsala University, Uppsala 75236, Sweden; and ²Department of Computer Science, Math & Science, American University of Paris, 31 Avenue Bosquet, Paris 75007, France

Summary

1. Maternal age effects on life-history traits, including longevity, are widespread and can be seen as a manifestation of ageing. However, little is known about how maternal life span may influence the maternal age effect. At a given chronological age, a long-lived parent may be at a younger biological age than a short-lived parent and thus has a less severe parental age effect. However, earlier work using experimentally evolved short- and long-lived lines did not support this hypothesis.
2. We scored developmental time and longevity of 14 995 individual seed beetles, *Callosobruchus maculatus* derived from replicate short-lived and long-lived lines created via artificial selection on male life span.
3. Offspring from older mothers had shorter life span, which is consistent with most of the literature.
4. We found support for the hypothesis that detrimental maternal age effects evolve to be weaker under selection for long life span. However, this finding was only apparent in males, suggesting that maternal age affects male and female offspring differently.
5. These results suggest that sex-dependent parental age effects should be incorporated in the studies of longevity and ageing evolution and that selection on one sex can cause evolution of parental age effects in the other sex.

Key-words: ageing, *Callosobruchus maculatus*, eclosion success, sex-specific response

Introduction

Ageing, defined as the decline in physiological and reproductive performance and the increase in probability of death with age, is a nearly universal phenomenon (Hamilton 1966; Rose 1991; Charlesworth 1994; Hughes & Reynolds 2005). Variation in the ageing rates, combined with different levels of extrinsic mortality, results in within and among species variation in life span. Life span is heritable (Johnson & Wood 1982; Klebanov *et al.* 2000; Fox *et al.* 2004a; Kemkes-Grottenthaler 2004), often sexually dimorphic (Trivers 1985; Bonduriansky *et al.* 2008; Maklakov & Lummaa 2013), and evolves rapidly in the laboratory (Rose 1984; Zwaan, Bijlsma & Hoekstra 1995; Partridge, Prowse & Pignatelli 1999; Berg & Maklakov 2012; Remolina *et al.* 2012).

Although life span is heritable, it can vary considerably among an individual's offspring. One key factor

contributing to such variation is parental age (Lansing 1947; Gavrilov & Gavrilova 1997; Priest, Mackowiak & Promislow 2002). The influence of parental age on offspring longevity has been a particularly hot topic in recent human studies, but has also been investigated in animal systems. The prevailing view is that offspring life span decreases with increased parental age (known as the 'Lansing effect'; Lansing 1947; Rockstein 1957; Tracey 1958; O'Brian 1961; Kiritani & Kimura 1967; Gavrilov & Gavrilova 1997; Priest, Mackowiak & Promislow 2002; García-Palomares *et al.* 2009; but see Fox, Bush & Wallin 2003). Moreover, the parental age effect is frequently specific to both parent and offspring sex. Maternal age is often the strongest factor affecting offspring life span (Butz & Hayden 1962; Priest, Mackowiak & Promislow 2002), although, intriguingly, in humans the father's age appears to play a bigger role (Gavrilov & Gavrilova 1997; Kemkes-Grottenthaler 2004).

A few studies have also investigated whether parental age affects sons and daughters differently. Interestingly, in humans, paternal age strongly influences the life span of daughters, while neither maternal nor paternal age had a

*Correspondence authors. E-mails: martin.i.lind@gmail.com; eberg@aup.edu

†These authors contributed equally to this study.

significant effect on the life span of sons (Gavrilov & Gavrilova 1997; Kemkes-Grottenthaler 2004). Unlike in humans, in model laboratory organisms, it has been possible to experimentally partition parental age effects and sex-specific offspring effects. However, there have been relatively few studies, and the results are mixed. One such study of fruit flies found that paternal age effects more strongly influenced the life span of sons, while maternal age effects more strongly influenced the life span of daughters (Priest, Mackowiak & Promislow 2002), the latter finding has also been reported in an early experiment by Butz & Hayden (1962). Also in mice, maternal age seems mostly to influence daughters (Carnes, Riesch & Schlupp 2012). In seed beetles, previous seminal work has shown that while maternal age affects males more than females, the overall effect was positive, contrary to a more common pattern observed in other systems (Fox, Bush & Wallin 2003).

Despite accumulating empirical evidence for parental age effects, very little is known about how parental life span should influence the parental age effect on offspring life span. This is important, since at a given chronological age, a long-lived parent may be at a younger biological age than a short-lived parent and thus has a less severe parental age effect. Since parental age effects are often caused by similar mechanisms that are involved in ageing (such as mutation load or trade-offs between early and late function) (Priest, Mackowiak & Promislow 2002; Kemkes-Grottenthaler 2004; Kong *et al.* 2012), it is likely that these effects will differ among long- and short-lived parents. Since the parental age effect has been shown to differ among genotypes (Priest, Mackowiak & Promislow 2002), it has the potential to be influenced by the evolution of parental life span.

We set out to test this hypothesis in long-lived and short-lived lines of the seed beetle *Callosobruchus maculatus* Fabricius (Berg & Maklakov 2012). The seed beetle *C. maculatus* is a model organism for studies of experimental life-history evolution (Messina 2004; Fricke & Arnqvist 2007; Maklakov, Bonduriansky & Brooks 2009), since it has a short generation time, thrives in a laboratory environment and is facultatively aphagous (i.e. does not require food or water once it emerges as an adult) (Fox 1993b; Fox, Bush & Wallin 2003). In this study, we used the lines that had been selected for long and short adult male life span, which resulted in the evolution of significant differences in longevity in both sexes because of intersexual genetic correlation for this trait (Berg & Maklakov 2012). We used these lines to test for sex- and selection-specific maternal age effects on offspring life span.

Materials and methods

STUDY SYSTEM

Callosobruchus maculatus is a very common pest of stored legumes. Females paste their eggs onto the host bean's surface.

Larvae hatch a few days later and burrow directly into the bean, using it as a food resource until hatching out as reproductively mature adults between 23 and 27 days after the egg is laid. *Callosobruchus maculatus* are capital breeders, obtaining all of the resources required for survival and reproduction during the larval stage (Fox 1993b; Fox, Bush & Wallin 2003). Females live shorter than males, and the adult life span is normally between 6 and 20 days, depending upon factors such as temperature and host plants (Fox, Czesak & Wallin 2004b; Fox *et al.* 2011; Berg & Maklakov 2012).

The long- and short-life selection lines used in our experiment were derived from a heterogeneous South Indian population ('SI USA') obtained from C. W. Fox at the University of Kentucky, USA. Originally collected in 1979 from infested mung beans (*Vigna radiata*) in Tirunelveli, India (Mitchell 1991), this stock population has been maintained in our laboratory for over 80 generations. The beetles have been cultured exclusively on mung beans and kept in climate chambers at 30 °C, 50% relative humidity and a 14:10 h light-dark cycle.

ARTIFICIAL SELECTION ON MALE LIFE SPAN

Prior to this experiment, we selected directly on male life span for a total of nine generations to create four replicate 'long-life' selection lines where males lived on average 40% longer than in four 'short-life' selection lines. For details of the selection procedure, see Berg & Maklakov (2012).

MATERNAL AGE EFFECTS

We allowed each of the lines to mate at random for two generations before assessing maternal age effects in order to reduce any residual parental effects. Beans, each bearing a single hatched egg, were then isolated in individual 'virgin chambers' (containers with separate wells for each bean) prior to hatching. For each of the four long-life and four short-life selection lines, a 1-day-old virgin female was paired with a 1-day-old virgin male from the unselected baseline population ($n = 20$ pairs per line), in order to eliminate any systematic male effects. Pairs were placed in a 60-mm Petri dish with *c.* 75 beans ('Day 1' dishes). We chose this number of beans, since females can lay up to 65 eggs per day (E. C. Berg, unpublished data), and we wanted to provide enough beans so that no more than one egg would be laid on each bean. After 24 h, the males were removed and discarded, females were moved to a new dish with 75 fresh beans ('Day 2' dishes) and the initial dishes were stored in the climate chamber. After 24 h, the female was moved to a new dish with 75 fresh beans ('Day 3'), and the Day 2 dishes were stored. This process was repeated one additional time ('Day 4+'). The females were allowed to remain in the fourth dish until death. Dishes were monitored daily, and the date of death of each female was recorded.

All hatched eggs from all days (Day 1 through Day 4+) were placed in individually marked virgin chambers and monitored daily until eclosion. The age at eclosion, sex and age at death were recorded for all offspring. This gives an accurate estimate of life span from all maternal ages, but because females were left in Day 4+, their eggs could be laid over several days, and therefore, the development time (but not life span) from Day 4+ will be overestimated. We therefore performed all analyses that included development time both with and without Day 4+.

We collected data on the offspring of 20 females of each of the eight lines. On average, we scored 94 offspring per female, resulting in a total of 14 995 offspring scored. Six thousand one hundred and forty-seven beetles emerged from eggs laid during Day 1, 2808 from Day 2, 2764 from Day 3 and 3236 from Day 4+.

STATISTICAL ANALYSIS

We tested the effect of maternal age and selection background on adult emergence success in a generalised mixed effect model implemented in a Bayesian MCMC framework using the package *MCMCglmm* (Hadfield 2010) in *R* 2.15.3 (R Development Core Team 2011). For all models, 1 day was subtracted from maternal age, to have the model intercept at Day 1. Emergence success was treated as a binary response variable (yes/no), selection background (long/short life) as a fixed factor and maternal age and age² as covariates estimating the direction (the age term) and curvature (the age² term) of the maternal age effect. Line was treated as a random effect, and because several offspring of each mother were scored, we also included maternal ID as a random effect to avoid pseudoreplication. Model selection using DIC (the Bayesian equivalent to AIC) was performed to find the minimal model.

All offspring that emerged from the beans were scored for development time and life span, which were investigated in separate mixed effect models with offspring sex and selection background as fixed factors, maternal age and age² as covariates and line and maternal ID were fitted as random effects. Response variables were log-transformed to meet the assumptions of normality. Since the maternal age effect influenced both development time and ageing, we also ran a similar model for life span that, in addition to the factors above, also included mean-centered development time as a covariate. Since development time from Day 4+ could be overestimated (see above), we analysed maternal age effects on development time, as well as the role of development time for life span both with and without the inclusion of Day 4+.

Results

Egg-to-adult survival was very high. On average, 97.6% of all eggs produced adults, and the eclosion success

increased significantly with increased maternal age (posterior mode: 0.366, 95% HPD interval: 0.127; 0.715, pMCMC: 0.015). The increase was linear, since the effect of maternal age² was not significant (posterior mode: -0.115, 95% HPD interval: -0.212; -0.004, pMCMC: 0.061). Selection background had no effect on egg-to-adult survival (posterior mode: 0.198, 95% HPD interval: -0.387; 0.563, pMCMC: 0.583).

Development time was affected by selection background, offspring sex and maternal age (Table 1, Fig. 1). We found that increased maternal age resulted in an increase in the development time of her offspring (with a positive quadratic component), but this effect was line specific. In lines selected for short male life span, offspring produced by young mothers took longer to develop than offspring from long-lived lines. However, this effect disappeared with increased maternal age (maternal age × selection treatment interaction), resulting in similar development time to the lines selected for long life span when laid by old mothers. Sons developed faster than daughters in both selection backgrounds, but the difference in development time between the selected lines was larger in daughters. A strong sign of a maternal age effect on development time in both selection backgrounds was also present when data from Day 4+ was included (Table 1).

We found that mothers from the lines selected for long life span produced offspring that had a longer life than offspring of mothers from short life span lines and that offspring life span in both selection backgrounds decreased with increased maternal age (Table 2, Fig. 2a). Daughters

Table 1. The influence of maternal age, selection treatment and offspring sex on the log of development time of the offspring. Results are presented separately for analyses including maternal age 1–3 and maternal age 1–4+

Parameter	Development time (maternal age 1–3)			Development time (maternal age 1–4+)		
	Posterior mode	95% HPD interval	pMCMC	Posterior mode	95% HPD interval	pMCMC
Intercept	3.127	3.103; 3.151	<0.001	3.129	3.106; 3.151	<0.001
Selection treatment (short → long)	-0.025	-0.059; 0.009	0.124	-0.026	-0.057; 0.008	0.117
Sex (female → male)	-0.019	-0.022; -0.017	<0.001	-0.019	-0.022; -0.017	<0.001
Maternal age	-0.007	-0.014; -0.001	0.041	-0.023	-0.027; -0.019	<0.001
Maternal age ²	0.004	0.001; 0.007	0.034	0.013	0.011; 0.014	<0.001
Selection treatment × Sex	0.004	0.001; 0.008	0.025	0.004	0.001; 0.008	0.024
Selection treatment × Maternal age	0.019	0.009; 0.027	<0.001	0.018	0.011; 0.022	<0.001
Selection treatment × Maternal age ²	-0.005	-0.009; 0.000	0.063	-0.003	-0.005; -0.001	<0.001




Fig. 1. Development time (\pm SE) in log days for (a) female (circles) and (b) male (triangles) offspring of mothers with an adult age of 1, 2, 3 or 4+ days from long-lived (closed symbols) or short-lived (open symbols) lines.

Table 2. The influence of maternal age, selection treatment and offspring sex on the log of offspring adult life span and life span residuals after controlling for development time. Life span is calculated on data from all maternal ages (1–4+), while life span residuals is calculated using data from the first three maternal ages

Parameter	Life span			Life span residuals		
	Posterior mode	95% HPD interval	pMCMC	Posterior mode	95% HPD interval	pMCMC
Intercept	3.294	3.244; 3.347	<0.001	3.307	3.254; 3.360	<0.001
Development time	–	–	–	–0.380	–0.434; –0.316	<0.001
Selection treatment (short → long)	0.050	–0.027; 0.118	0.217	0.032	–0.043; 0.110	0.346
Sex (female → male)	–0.367	–0.380; –0.355	<0.001	–0.378	–0.391; –0.366	<0.001
Maternal age	–0.036	–0.054; –0.018	<0.001	–0.087	–0.109; –0.059	<0.001
Maternal age ²	0.007	0.002; 0.014	0.020	0.035	0.021; 0.046	<0.001
Selection treatment × Sex	0.043	0.024; 0.057	<0.001	0.048	0.030; 0.064	<0.001
Selection treatment × Maternal age	–0.021	–0.046; 0.003	0.086	0.043	0.014; 0.072	0.004
Selection treatment × Maternal age ²	0.005	–0.002; 0.014	0.136	–0.028	–0.042; –0.013	<0.001
Sex × Maternal age	–0.084	–0.102; –0.053	<0.001	–0.070	–0.102; –0.042	<0.001
Sex × Maternal age ²	0.029	0.020; 0.037	<0.001	0.025	0.009; 0.038	0.002
Selection treatment × Sex × Maternal age	0.088	0.062; 0.130	<0.001	0.040	0.026; 0.056	<0.001
Selection treatment × Sex × Maternal age ²	–0.031	–0.044; –0.021	<0.001	–	–	–




Fig. 2. (a) Adult life span and (b) residuals of adult life span after controlling for the effect of development time in log days (\pm SE) for female (circles) and male (triangles) offspring of mothers with an adult age of 1, 2, 3 or 4+ days from long-lived (closed symbols) or short-lived (open symbols) lines. Life span residuals were converted to original units by adding the grand mean.

lived substantially longer than sons, and the maternal age effect was specific to both offspring sex and the selection background of the line. The maternal age effect was stronger for sons from short-life lines than for sons from long-life lines, but this effect was most pronounced in offspring born during the first 2 days of a mother's adult life span, as indicated by the interaction between selection treatment, sex and maternal age. We also found that the quadratic curvature differed between lines and offspring sex, most

noticeable for males of the short-lived lines, who not only had the largest drop in offspring life span between maternal age 1 and 2, but also a small increase in offspring life span for age 3 and 4+. No obvious difference in maternal age effect between the selected lines was present for daughters.

When development time was used as a covariate, we found that long development time was associated with short life span, with a similar effect in both sexes and

selection treatments (Table 2, Fig. 2b). Much of the difference in life span between the lines was explained by development time, but not the sex difference in life span. Offspring born to young (Day 1) mothers had longer life span than expected when controlling for development time, suggesting that other factors than development time are responsible for the maternal age effect on offspring life span. Again, an interaction between selection treatment, sex and maternal age indicated that the negative effect on maternal age in the short-lived lines was most pronounced in sons produced during the first 2 days. An interaction between offspring sex and maternal age² indicates a different shape of the maternal age effect between sons and daughters, and this effect was present in all selected lines. When data from the Day 4+ treatment was included, it did not affect the pattern found (see Table S1, Supporting information).

Discussion

We found that offspring of older mothers had reduced life span, which is consistent with the large body of literature on the negative impact of older parents across the animal kingdom (Lansing 1947; Rockstein 1957; Tracey 1958; O'Brian 1961; Butz & Hayden 1962; Kiritani & Kimura 1967; Klass 1977; GavriloV & GavriloV 1997; Priest, Mackowiak & Promislow 2002; Kemkes-Grotenthaler 2004; Tucić, Šešlija & Stanković 2004; but see Fox, Bush & Wallin 2003). Our main finding, however, is that the maternal age effect on offspring life span had evolved in long-lived lines and that this evolutionary response was sex-specific: increased maternal age is less damaging for offspring life span when mothers are from a long-lived genetic background, and this effect is stronger for sons than for daughters. This effect was, however, only present during the first 2 days of maternal age. We found that development time increased with maternal age, likely because of a reduction in maternal provisioning, but that the sex-specific maternal age effect on offspring life span was present even after controlling for the strong negative effect of long development time on life span.

Maternal age effects that reduce offspring life span are common (but see Fox, Bush & Wallin 2003; discussed below) and are believed to be caused by a number of mechanisms including maternal provisioning, the accumulation of late acting deleterious mutations and trade-offs between early- and late-life function (Priest, Mackowiak & Promislow 2002). Since these two latter mechanisms are also key explanations for the evolution of life span and ageing and that genetic variation for maternal age effects is present in nature (Priest, Mackowiak & Promislow 2002), there is a potential for maternal age effects to be influenced by the evolution of ageing. We therefore investigated whether lines artificially selected for long or short male life span would also differ in maternal age effects. We found not only that individuals whose mothers were from lines selected for long life lived longer, but also that the

maternal age effect that caused a reduction in offspring life span was weaker in the long-life lines, especially for sons. Since long-lived females likely had a lower biological age at a given chronological age compared to the short-lived females, this result suggests that long maternal life span reduces the negative effect of maternal age on offspring life span. Notably, the effect was most pronounced during the first days of female adult life, with a large drop in offspring life span laid by 2-day-old mothers from the short-lived lines. Female *C. maculatus* are known to lay most eggs during the first days of adult life (Fox 1993a; Tatar & Carey 1995), indeed 60% of total eggs produced in this experiment were laid during the first 2 days. This suggests that the maternal age effect may have a large fitness impact.

Interestingly, the only previous study that investigated the evolution of maternal age effects in short-lived and long-lived experimental lines reported negative results. When comparing lines of the bean weevil *Acanthoscelides obtectus* selected for early or late reproduction, no difference in the maternal age effect on offspring life span between the selected lines was found (Tucić, Šešlija & Stanković 2004). The difference between the studies may lie in the biology of these species suggesting that evolution of long life may not always manifest itself in the reduction of detrimental parental age effects. Clearly, more experimental work is needed before we can make broad cross-taxonomic generalizations regarding the evolution of parental age effects in response to selection for age-specific life-histories.

The evolution of maternal age effect in our study was specific to offspring sex: the reduction of the maternal age effect in long-lived lines mainly affected sons, not daughters. While sex difference in maternal age effects on offspring life span is common, there is substantial variation across species, and even populations within species, regarding which sex is affected stronger (Butz & Hayden 1962; Priest, Mackowiak & Promislow 2002; Fox, Bush & Wallin 2003; Carnes, Riesch & Schlupp 2012). Generally, our finding of stronger response in males is in line with stronger maternal effects in general (Fox, Czesak & Wallin 2004b) and maternal age effects in particular (Fox, Bush & Wallin 2003) on male life span in *C. maculatus*. Sexual selection often causes males to invest heavily into energetically costly traits (Sheldon *et al.* 1998; Brooks 2000; Tomkins *et al.* 2004; Bussiere *et al.* 2008; Kwan *et al.* 2008; Sharp & Agrawal 2013), and recent work found that male *C. maculatus* are indeed more sensitive to environmental changes (Berger *et al.* 2014). Male *C. maculatus* spend a lot of energy in mate search even in the absence of females; therefore, low-condition sons produced by older mothers could be paying relatively higher price than low-condition daughters.

Since a well-known maternal age effect in *C. maculatus* is a reduction in maternal investment in egg size, resulting in longer development time for offspring hatching from smaller eggs (Fox 1993b, 1994), we also investigated the maternal age effect on development time. In agreement

with previous work, we found that offspring from older mothers took longer to develop. In addition, we found a strong effect of our selection treatment, where offspring from long-life lines had shorter development times, a difference that was especially pronounced for young mothers, while the maternal age effect on development time was weak in the short-lived lines. Although we did not measure egg size, the pattern is consistent with previous findings in *C. maculatus* of a negative correlation between egg size and development time, and reduced egg size and increased offspring development time with maternal age (Fox 1993b, 1994). Since the females from long-life lines were larger than females from short-life lines (Berg *et al.* submitted), and there is a positive correlation between female body size and egg size in *C. maculatus* (Fox 1993b), our results suggest that the large females of the long-life lines lay large eggs, especially when young, resulting in offspring with short development time and long life span. Although neither egg size nor offspring size was measured here, previous research suggests there is no maternal age effect on offspring size in *C. maculatus* (Fox 1993b), despite the fact that old mothers lay small eggs, possibly because they take longer to develop and still hatch at a common size.

Although the maternal age effect on development time, probably caused by a reduction in egg size with age, is a likely mechanism explaining a major part of the parental age effect, this effect alone does not explain the patterns we found. By investigating the maternal age effect on life-span after controlling for the strongly negative effect of increased development time, the maternal age effect was still present, resulting in longer life span than expected for offspring born to newly eclosed females. Not surprisingly, the life spans of beetles from the two selection lines were more similar after controlling for development time, since the lines selected for long life had shorter development time. It is possible that ageing of eggs caused this effect or other factors related to maternal provisioning that do not affect development time. It is also interesting to note that also Fox *et al.* (2011) have found sex-specific responses in life span, but in their case as a response to novel conditions, further suggesting that the life span of the two sexes respond differently in *C. maculatus*.

While our study is consistent with most of the literature on parental age effects that show that older parents produce shorter-lived offspring, our results differ from the only other similar study on *C. maculatus*, in which Fox, Bush & Wallin (2003) found a positive effect of maternal age on offspring life span, but only for very old mothers (Day 6+) which are outside the range of our study. The authors hypothesized that their unexpected results may have been caused by the reduced eclosion success of offspring from older mothers in their experiment, resulting in condition-dependent survival. In other words, the subset of offspring from old mothers that did eclose were in high condition and, therefore, lived longer. In our study, we did not find a decrease in eclosion success with maternal age (rather a small increase) and, consequently, condition-dependent mortality did not

influence our estimate of the maternal age effect. The fact that our results showed negative effects of maternal age suggest that differences in egg-to-adult survival, as argued by Fox, Bush & Wallin (2003), can influence the estimates of the maternal age effect on life span.

Selection on late age of reproduction (Luckinbill *et al.* 1984; Rose 1984; Partridge, Prowse & Pignatelli 1999) or direct selection on long-lived individuals (Zwaan, Bijlsma & Hoekstra 1995; Hunt *et al.* 2006; Berg & Maklakov 2012) result in the evolution of long life, but at the same time the offspring of old parents have reduced life span (Lansing 1947; Priest, Mackowiak & Promislow 2002). Therefore, the processes driving maternal age effects have often been seen as distinct from those underlying ageing. Here, we show that the detrimental effect of old maternal age on the life span of offspring evolves rapidly and is diminished in mothers from long-lived lines. Moreover, sons were affected more than daughters, suggesting that maternal age effects can play an important role in shaping sexual dimorphism in life span and ageing. More broadly, our findings confirm that parental age effects can be viewed as a manifestation of ageing (Priest, Mackowiak & Promislow 2002) and call for a further integration of the two fields.

Acknowledgements

The study was supported by the Swedish Research Council and the ERC Starting Grant 2010 to A.A.M., the Carl Tryggers Stiftelse and Zoologiska Stiftelsen to E.C.B and the Swedish Research Council to M.I.L. We thank Cindy Canton for help with data collection in the laboratory, and the editor, associate editor and two anonymous reviewers for constructive comments on a previous version of this manuscript.

Data accessibility

Data deposited in the Dryad repository: <http://doi.org/10.5061/dryad.2320n> (Lind *et al.* 2015).

References

- Berg, E.C. & Maklakov, A.A. (2012) Sexes suffer from suboptimal lifespan because of genetic conflict in a seed beetle. *Proceedings of the Royal Society of London. Series B, Biological Sciences*, **279**, 4296–4302.
- Berg, E.C., Lind, M.I., Alavioon, G. & Maklakov, A.A. (2014) Data from: evolution of differential maternal age effects on male and female offspring development and longevity. *Dryad Digital Repository*, doi:10.5061/dryad.2320n.
- Berger, D., Grieshop, K., Lind, M.I., Goenaga, J., Maklakov, A.A. & Arnqvist, G. (2014) Intralocus sexual conflict and environmental stress. *Evolution*, Early View, doi:10.1111/evo.12439.
- Bonduriansky, R., Maklakov, A., Zajitschek, F. & Brooks, R. (2008) Sexual selection, sexual conflict and the evolution of ageing and life span. *Functional Ecology*, **22**, 443–453.
- Brooks, R. (2000) Negative genetic correlation between male sexual attractiveness and survival. *Nature*, **406**, 67–70.
- Bussiere, L.F., Hunt, J., Stölting, K.N., Jennions, M.D. & Brooks, R. (2008) Mate choice for genetic quality when environments vary: suggestions for empirical progress. *Genetica*, **134**, 69–78.
- Butz, A. & Hayden, P. (1962) The effects of age of male and female parents on the life cycle of *Drosophila melanogaster*. *Annals of the Entomological Society of America*, **55**, 617–618.
- Carnes, B.A., Riesch, R. & Schlupp, I. (2012) The delayed impact of parental age on offspring mortality in mice. *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences*, **67A**, 351–357.

- Charlesworth, B. (1994) *Evolution in Age-Structured Populations*, 2nd edn. Cambridge University Press, New York, NY, USA.
- Fox, C.W. (1993a) Multiple mating, lifetime fecundity and female mortality of the bruchid beetle *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Functional Ecology*, **7**, 203–208.
- Fox, C.W. (1993b) The influence of maternal age and mating frequency on egg size and offspring performance in *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Oecologia*, **96**, 139–146.
- Fox, C.W. (1994) The influence of egg size on offspring performance in the seed beetle, *Callosobruchus maculatus*. *Oikos*, **71**, 321–325.
- Fox, C.W., Bush, M.L. & Wallin, W.G. (2003) Maternal age affects offspring lifespan of the seed beetle, *Callosobruchus maculatus*. *Functional Ecology*, **17**, 811–820.
- Fox, C.W., Czesak, M.E. & Wallin, W.G. (2004b) Complex genetic architecture of population differences in adult lifespan of a beetle: nonadditive inheritance, gender differences, body size and a large maternal effect. *Journal of Evolutionary Biology*, **17**, 1007–1017.
- Fox, C.W., Bush, M.L., Roff, D.A. & Wallin, W.G. (2004a) Evolutionary genetics of lifespan and mortality rates in two populations of the seed beetle, *Callosobruchus maculatus*. *Heredity*, **92**, 170–181.
- Fox, C.W., Wagner, J.D., Cline, S., Thomas, F.A. & Messina, F.J. (2011) Rapid evolution of lifespan in a novel environment: sex-specific responses and underlying genetic architecture. *Evolutionary Biology*, **38**, 182–196.
- Fricke, C. & Arnqvist, G. (2007) Rapid adaptation to a novel host in a seed beetle (*Callosobruchus maculatus*): the role of sexual selection. *Evolution*, **61**, 440–454.
- García-Palomares, S., Navarro, S., Pertusa, J.F., Hermenegildo, C., García-Pérez, M.A., Rausell, F. et al. (2009) Delayed fatherhood in mice decreases reproductive fitness and longevity of offspring. *Biology of Reproduction*, **80**, 343–349.
- Gavrilov, L.A. & Gavrilova, N.S. (1997) Parental age at conception and offspring longevity. *Reviews in Clinical Gerontology*, **7**, 5–12.
- Hadfield, J. (2010) MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of Statistical Software*, **33**, 1–22.
- Hamilton, W.D. (1966) The moulding of senescence by natural selection. *Journal of Theoretical Biology*, **12**, 12–45.
- Hughes, K.A. & Reynolds, R.M. (2005) Evolutionary and mechanistic theories of aging. *Annual Review of Entomology*, **50**, 421–445.
- Hunt, J., Jennions, M.D., Spyrou, N. & Brooks, R. (2006) Artificial selection on male longevity influences age-dependent reproductive effort in the black field cricket *Telegryllus commodus*. *The American Naturalist*, **168**, E72–E86.
- Johnson, T.E. & Wood, W.B. (1982) Genetic analysis of life-span in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences*, **79**, 6603–6607.
- Kemkes-Grottenthaler, A. (2004) Parental effects on offspring longevity—evidence from 17th to 19th century reproductive histories. *Annals of Human Biology*, **31**, 139–158.
- Kiritani, K. & Kimura, K. (1967) Effects of parental age on the life cycle of the southern green stink bug, *Nezara viridula*. *Applied Entomology and Zoology*, **2**, 69–78.
- Klass, M.R. (1977) Aging in the nematode *Caenorhabditis elegans*: major biological and environmental factors influencing life span. *Mechanisms of Ageing and Development*, **6**, 413–429.
- Klebanov, S., Flurkey, K., Roderick, T.H., Archer, J.R., Astle, M.C., Chen, J. et al. (2000) Heritability of life span in mice and its implication for direct and indirect selection for longevity. *Genetica*, **110**, 209–218.
- Kong, A., Frigge, M.L., Masson, G., Besenbacher, S., Sulem, P., Magnusson, G. et al. (2012) Rate of *de novo* mutations and the importance of father's age to disease risk. *Nature*, **488**, 471–475.
- Kwan, L., Bedhomme, S., Prasad, N.G. & Chippindale, A.K. (2008) Sexual conflict and environmental change: trade-offs within and between the sexes during the evolution of desiccation resistance. *Journal of Genetics*, **87**, 383–394.
- Lansing, A.I. (1947) A transmissible, cumulative, and reversible factor in aging. *Journal of Gerontology*, **2**, 228–239.
- Lind, M.I., Berg, E.C., Alavioion, G. & Maklakov, A.A. (2015) Data from: Evolution of differential maternal age effects on male and female offspring development and longevity. *Dryad Digital Repository*, <http://doi.org/10.5061/dryad.2320n>.
- Luckinbill, L.S., Arking, R., Clare, M.J., Cirocco, W.C. & Buck, S.A. (1984) Selection for delayed senescence in *Drosophila melanogaster*. *Evolution*, **38**, 996–1003.
- Maklakov, A.A., Bonduriansky, R. & Brooks, R.C. (2009) Sex differences, sexual selection, and ageing: an experimental evolution approach. *Evolution*, **63**, 2491–2503.
- Maklakov, A.A. & Lummaa, V. (2013) Evolution of sex differences in life-span and ageing: causes and constraints. *BioEssays*, **35**, 717–724.
- Messina, F.J. (2004) Predictable modification of body size and competitive ability following a host shift by a seed beetle. *Evolution*, **58**, 2788–2797.
- Mitchell, R. (1991) The traits of a biotype of *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) from South India. *Journal of Stored Products Research*, **27**, 221–224.
- O'Brian, D.M. (1961) Effects of parental age on the life cycle of *Drosophila melanogaster*. *Annals of the Entomological Society of America*, **54**, 412–416.
- Partridge, L., Prowse, N. & Pignatelli, P. (1999) Another set of responses and correlated responses to selection on age at reproduction in *Drosophila melanogaster*. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **266**, 255–261.
- Priest, N.K., Mackowiak, B. & Promislow, D.E.L. (2002) The role of parental age effects on the evolution of aging. *Evolution*, **56**, 927–935.
- R Development Core Team. (2011) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Remolina, S.C., Chang, P.L., Leips, J., Nuzhdin, S.V. & Hughes, K.A. (2012) Genomic basis of aging and life-history evolution in *Drosophila melanogaster*. *Evolution*, **66**, 3390–3403.
- Rockstein, M. (1957) Longevity of male and female house flies. *Journal of Gerontology*, **12**, 253–256.
- Rose, M.R. (1984) Laboratory evolution of postponed senescence in *Drosophila melanogaster*. *Evolution*, **38**, 1004–1010.
- Rose, M.R. (1991) *Evolutionary Biology of Aging*. Oxford University Press, New York, NY.
- Sharp, N.P. & Agrawal, A.F. (2013) Male-biased fitness effects of spontaneous mutations in *Drosophila melanogaster*. *Evolution*, **67**, 1189–1195.
- Sheldon, B.C., Merilä, J., Lindgren, G. & Ellegren, H. (1998) Gender and environmental sensitivity in nestling collared flycatchers. *Ecology*, **79**, 1939–1948.
- Tatar, M. & Carey, J.R. (1995) Nutrition mediates reproductive trade-offs with age-specific mortality in the beetle *Callosobruchus maculatus*. *Ecology*, **76**, 2066–2073.
- Tomkins, J.L., Radwan, J., Kotiaho, J.S. & Tregenza, T. (2004) Genic capture and resolving the lek paradox. *Trends in Ecology & Evolution*, **19**, 323–328.
- Tracey, K.M. (1958) Effects of parental age on the life cycle of the mealworm, *Tenebrio molitor* Linnaeus. *Annals of the Entomological Society of America*, **51**, 429–432.
- Trivers, R. (1985) *Social Evolution*. Benjamin/Cummings Publishing Company, Inc., Menlo Park, CA, USA.
- Tucić, N., Šešljica, D. & Stanković, V. (2004) The short-term and long-term effects of parental age in the bean weevil (*Acanthoscelides obtectus*). *Evolutionary Ecology*, **18**, 187–201.
- Zwaan, B., Bijlsma, R. & Hoekstra, R.F. (1995) Direct selection on life span in *Drosophila melanogaster*. *Evolution*, **49**, 649–659.

Received 7 November 2013; accepted 13 June 2014
Handling Editor: Wolf Blanckenhorn

Supporting Information

Additional Supporting information may be found in the online version of this article:

Table S1. The influence of maternal age (Day 1–4+), selection treatment and offspring sex on the log of offspring life span, after controlling for offspring development time.